


- Anisomeridium americanum* (A. Massal.) R.C. Harris, More Florida Lichens, Incl. 10 Cent Tour Pyrenol. (New York): 144 (1995)  
 = *Arthopyrenia americana* A. Massal., Ric. auton. lich. crost. (Verona): 170, fig. 341 (1852)  
 = *Arthopyrenia limitans* Müll. Arg., Flora, Regensburg 66(19): 306 (1883)  
 = *Ditremis americana* (A. Massal.) R.C. Harris, Some Florida Lichens (New York): 31 (1990)  
 = *Ditremis pyrenuloides* (Müll. Arg.) Makhija & Patw., Biovigyanam 16(1): 22 (1990)  
 = *Leiophloea americana* (A. Massal.) Trevis., Conspect. Verruc.: 9 (1860)  
 = *Pleurotrema pyrenuloides* Müll. Arg., Rep. Meetings Australs. Assoc. Advancem. Sci.: 462 (1895)  
 = *Pyrenula americana* (A. Massal.) Trevis., Spighe Paglie: 18 (1853)



Thallus endophloeodal, not corticate, whitish to greyish, UV-. Ostiolum excentric. Ascospores 8/ascus, hyaline, fusiform, fine granular ornamented, 1-septate, with septum  $\pm$  submedian, 26-35(-42) x 7.5-10(-12)  $\mu$ m

[ABL18829], Brazil, Sergipe, Santa Luzia do Itanhy, Mata do Crasto, 11°22' S, 37°25' W, 10 m. On tree bark in Atlantic rainforest. Leg. A. Aptroot & M. Cáceres (no 18829), 16.03.2014, det. A. Aptroot, 2014.







*Anisomeridium americanum*



*Anisomeridium americanum*



*Anisomeridium americanum*