


- Anisomeridium terminatum* (Nyl.) R.C. Harris [as 'terminata'], More Florida Lichens, Incl. 10 Cent Tour Pyrenol. (New York): 130 (1995)
= *Arthopyrenia terminata* (Nyl.) Müll. Arg., Flora, Regensburg 73: 345 (1890)
= *Ditremis terminata* (Nyl.) R.C. Harris, in Egan, Bryologist 90(2): 163 (1987)
= *Tomasellia queenslandica* Müll. Arg., Rep. Meetings Australs. Assoc. Advancem. Sci.: 460 (1895)
= *Verrucaria terminata* Nyl., Bull. Soc. linn. Normandie, sér. 2 7: 180 (1873)

Thallus UV-. Ostiole lateral. Ascospores 8/ascus, hyaline, 1-septate, 18-27 x 5.5-7.5 μm

[ISE40357], Brazil, Pará, Fazenda Pantera, 85 km N of Dom Eliseu, in primary tropical lowland rain forest on bark of tree, 3°49'54" S, 48°03'37" W, 120 m. Leg. M. Cáceres & A. Aptroot (no 40357), 29.10.2016, det. A. Aptroot, 3016.







Anisomeridium terminatum



Anisomeridium terminatum



Anisomeridium terminatum