

- Letrouitia domingensis* (Pers.) Hafellner & Bellem., Nova Hedwigia 35(2 & 3): 281 (1982) [1981]  
 = *Bombyliospora domingensis* (Pers.) Zahlbr., in Wawra & Beck, Itin. Princ. S. Coburgi 2: 153 (1888)  
 = *Heterothecium domingense* (Pers.) Tuck., Proc. Amer. Acad. Arts & Sci. 7: 229 (1868) [1866]  
 = *Lecanora domingensis* (Pers.) Ach., Syn. meth. lich. (Lund): 336 (1814)  
 = *Lecidea domingensis* (Pers.) Nyl., Acta Soc. Sci. fenn. 7(2): 462 (1863)  
 = *Lopadium domingense* (Pers.) Fink, Lich. Fl. U.S.: 235 (1935)  
 = *Miltidea domingensis* (Pers.) Stirt., Trans. Proc. N.Z. Inst. 30: 386 (1898) [1897]  
 = *Parmelia ventosa* var. *domingensis* (Pers.) Eschw., in Martius, Fl. bras. enum. pl. 1(1): 189 (1833)  
 = *Patellaria domingensis* Pers., Ann. Wetter. Gesellsch. Ges. Naturk. 2(1): 12 (1810)  
 = *Placodium domingense* (Pers.) Vain., Dansk bot. Ark. 4(no. 11): 11 (1926)

[ABL48980], Brazil, Sena Madureira, Reserva extratevista Cazumbá-Iracema, Comunidade Cuidado, on tree bark in disturbed tropical rain forest. 09°07' S, 68°59' W, 150 m. Leg. A. Aptroot (no 48980), M. Cáceres & J. Cavalcante, 09.04.2019. Det. A. Aptroot. 2019.





*Letrouitia domingensis*