

- Letrouitia subvulpina*** (Nyl.) Hafellner, *Nova Hedwigia* 35(4): 705 (1983)  
 [1981]  
 = *Heterothecium sayeri* Müll. Arg., *Flora, Regensburg* 70: 338 (1887)  
 = *Lecidea sayeri* (Müll. Arg.) Shirley, *Proc. R. Soc. Qd.* 6: 184 (1889)  
 = *Lecidea subvulpina* Nyl., *Bull. Soc. linn. Normandie, sér. 2* 2: 89 (1868)  
 = *Letrouitia sayeri* (Müll. Arg.) Elix, *Australas. Lichenol.* 61: 23 (2007)  
 = *Lopadium sayeri* (Müll. Arg.) Zahlbr., *Cat. Lich. Univers.* 4: 313 (1926)  
 [1927]  
 = *Lopadium vulpinum* var. *subvulpinum* (Nyl.) Zahlbr., *Cat. Lich. Univers.* 4: 317 (1926) [1927] *Nat. Pflanzenfam., Teil. I* (Leipzig) 1(1\*): 137 (1905)

[ABL50207], Brazil, Sena Madureira, Reserva extratevista Cazumbá-Iracema, Núcleo Cazumbá, on tree bark in disturbed tropical rain forest. 09°07' S, 68°57' W, 150 m. Leg. A. Aptroot (no 50207), M. Cáceres & J. Cavalcante, 10.04.2019. Det. A. Aptroot. 2019.





*Letrouitia subvulpina*