Myriotrema glaucophaenum (Kremp.) Hale
[= ***Myriotrema subcostaricense*** Sipman]

Thallus corticolous, grey-olive, continuous, verruculose to finely warty. Apothecia moderately emergent to sessile, rounded to angular, 0.5-1.1 mm wide, with a open and fissured pore. Proper exciple fused or the tips incompletely free. Columella absent. Disc covered by thin white pruina. Ascospores hyaline, transversely septate, 4-6 loculate, 13-24 x 5-8 µm, J+ blue-violet. Chemistry: psoromic acid (maj). Pantropical

[17225], Venezuela/Aragua. Abfahrt auf der Straße von Ocumare de la Costa nach Maracay, unterhalb der biologischen Station "Dr. Alberto Fernandez Y.". An einem Laubbaum in einem stark gestörten , verbuschten Regenwald. 200 m. 10°22'32" N, 67°43'10" W. Leg. K. Kalb & J. Hernandez 05.08.2010, det. K. Kalb. Aus Kalb: Lichenes Neotropici 593.





Myriotrema glaucophaenum





Myriotrema glaucophaenum





Myriotrema glaucophaenum



Myriotrema glaucophaenum

10 µm

J+ violet

10 µm



10 µm



20 µm

thallus

Myriotrema glaucophaenum