





Astrothelium subdiscretum (Nyl.) Aptroot & Lücking, Lichenologist 48(6): 890 (2016)
= *Bathelium subdiscretum* (Nyl.) Müll. Arg., Bot. Jb. 6: 395 (1885)
= *Laurera subdiscreta* (Nyl.) Zahlbr., Cat. Lich. Univers. 1: 506 (1922)
= *Trypethelium subdiscretum* Nyl., Flora, Regensburg 52: 73 (1869)
= *Laurera fusispora* Makhija & Patw., Mycotaxon 31:576 (1988).

Thallus corticate, olive-green to brownish, uneven to verrucose-rugulose. Ascomata trypethelioid, with apical ostioles, pseudostromatic; pseudostromata 0.8-1.5(-2.0) mm broad, immersed-erumpent, irregular to linear and often confluent and reticulate, with ascomata exposed and black. Hamathecium interspersed or clear. Asospores 8/ascus, 13-17-septate, 35-50 x 12-18 μm , without distinctly thickened median septum, hyaline, IKI-. Chemistry: Thallus and pseudostromata UV-, K-. TLC: no substances detected.

[AMH742499], India, Karnataka, Anamod ghat, Anamod-Goa Road. Leg. Patwardhan (no AMH7424399) & Prabhu, 10.12.1974. ISOTYPE of *Laurera fusispora* Makhija & Patw.





Astrothelium subdiscretum