

- Dictyomeridium amylosporum* (Vain.) Aptroot, M.P. Nelsen & Lücking, in Aptroot & Lücking, Lichenologist 48(6): 923 (2016)  
= *Campylothelium amylosporum* (Vain.) R.C. Harris, in Tucker & Harris, Bryologist 83(1): 7 (1980)  
= *Polyblastiopsis amylospora* (Vain.) Zahlbr., Cat. Lich. Univers. 1: 347 (1922)  
= *Polymeridium amylosporum* (Vain.) Aptroot, Nova Hedwigia 98(1-2): 10 (2013) [2014]  
= *Thelenella amylospora* Vain., Acta Soc. Fauna Flora fenn. 7(no. 2): 218 (1890)

Thallus ecorticate, white, UV-. Ascomata pleurothelioid, with lateral ostioles, solitary, 0.4-0.6 mm diam., erumpent to prominent, black. Hamathecium clear, filaments profusely anastomosing. Ascospores 8/ascus, IKI+ violet, muriform, 44-55 x 12-21  $\mu\text{m}$ , with partly oblique septa, not ornamented, wall not thickened. Chemistry: Thallus and ascomata UV-, K-. TLC: no substances detected.

[ABL41150], Brazil, Minas Gerais, Catas Atlas, Serra do Caraça, Parque Nacional do Caraça, near Banho do Belchior, 20°06' S, 43°20' W, 1250 m. Leg. A. Aptroot (no 41150), 17.09.1997, det. A. Aptroot, 1998.







*Dictyomeridium amylosporum*