

- Dictyomeridium proponens*** (Nyl.) Aptroot, M.P. Nelsen & Lücking, in Lücking et al., *Lichenologist* 48(6): 757 (2016)  
 = *Campylothelium proponens* (Nyl.) Müll. Arg., *Hedwigia* 31: 286 (1892)  
 = *Polyblastia proponens* (Nyl.) Müll. Arg., *Flora, Regensburg* 65: 402 (1882)  
 = *Polyblastiopsis proponens* (Nyl.) Zahlbr., *Cat. Lich. Univers.* 1: 351 (1922)  
 = *Polymeridium proponens* (Nyl.) R.C. Harris, *Bolm Mus. paraense 'Emílio Goeldi', sér. bot.* 7(2): 637 (1993) [1991]  
 = *Verrucaria proponens* Nyl., *Bull. Soc. linn. Normandie, sér. 2* 2: 130 (1868)

Thallus ecorticate, white. Ascomata pleurothelioid, with lateral ostioles, solitary, 0.4-0.6 mm diam., erumpent, grey black. Hamathecium clear, filaments profusely anastomosing. Ascospores 8/ascus, IJI+ violet, muriform, 33-54 x 12-19  $\mu\text{m}$ , with partly oblique septa, not ornamented, wall not thickened. Chemistry: Thallus UV+ yellow, K-. TLC: lichexanthone.

[ABL41223], Brazil, Minas Gerais, Catas Atlas, Serra do Caraça, Parque Natural do Caraça near Funil, 1 km NW of monastery Santuário do Caraça, in *Eremanthus* shrub on tree. 20°06' S, 43°29' W c. 1250 m. Leg. A. Aptroot (no 41223), 18.09.1997, det. A. Aptroot 1998.





*Dictyomeridium proponens*