


- Trinathotrema stictideum* (Nyl.) Lücking, R. Miranda & Kalb, in Lücking, Rivas Plata, Mangold, Sipman, Aptroot, Miranda González, Kalb, Chaves, Ventura & Esqui, Biblthca Lichenol. 106: 200 (2011)**
- = *Conotrema albonigrum* Zahlbr., Annls mycol. 33: 42 (1935)
  - = *Conotrema stictideum* (Nyl.) G. Salisb., Nova Hedwigia 25(3-4): 691 (1974)
  - = *Gyalecta stictidea* (Nyl.) Zahlbr., Cat. Lich. Univers. 2: 714 (1924)
  - = *Lecidea stictidea* Nyl., Flora, Regensburg 49: 290 (1866)
  - = *Ocellularia gyrostromoides* Müll. Arg., Flora, Regensburg 71: 47 (1888)
  - = *Ocellularia scolecospora* C.W. Dodge, Ann. Mo. bot. Gdn 40: 345 (1953)
  - = *Ocellularia stictidea* (Nyl.) Vězda, Ark. Bot., Ser. 2 7: 177 (1969)
  - = *Ocellularia texana* (Willey) Zahlbr., Cat. Lich. Univers. 2: 602 (1923) [1924]
  - = *Schizoxylon gyrostromoides* (Müll. Arg.) Lumbsch & Papong, Mycota-xon 109: 320 (2009)
  - = *Thelotrema texanum* Willey, Acta Soc. Sci. fenn. 26(no. 10): 17 (1900)

[ABL50138], Brazil, Sena Madureira, Reserva extratevista Cazumbá-Iracema, near Colonia Esperanza, on tree bark in primary tropical rain forest. 09°06' S, 68°58' W, 140 m. Leg. A. Aptroot (no 50138), M. Cáceres & J. Cavalcante, 10.04.2019. Det. A. Aptroot. 2019.





*Trinathotrema stictideum*